

Fuzzy Anti-bounded Linear Operator

Bivas Dinda, T.K. Samanta and Iqbal H. Jebril

Department of Mathematics,
Mahishamuri Ramkrishna Vidyapith, West Bengal, India.

e-mail: bvsdinda@gmail.com

Department of Mathematics, Uluberia College, West Bengal, India.

e-mail: mumpu_tapas5@yahoo.co.in

Department of Mathematics, King Faisal University, Saudi Arabia.

e-mail: iqbal501@hotmail.com

Abstract

Various types of fuzzy anti-continuity and fuzzy anti-boundedness are defined. A few properties of them are established. The intra and inter relation among various types of fuzzy anti-continuity and fuzzy anti-boundedness are studied.

Keywords: *Fuzzy anti-norm, α -norm, Fuzzy α -anti-convergence, Fuzzy anti-continuity, Fuzzy anti-boundedness.*

2010 Mathematics Subject Classification: 03E72, 46S40.

1 Introduction

Fuzzy set theory was first introduced by Zadeh [13] in 1965 and thereafter several authors contributed different articles on this concept and applied on different branches of pure and applied mathematics. The concept of fuzzy norm was introduced by Katsaras [9] in 1984. In 1992, Felbin [7] introduced the idea of fuzzy norm on a linear space. Cheng-Moderson [4] introduced another idea of fuzzy norm on a linear space whose associated metric is same as the associated metric of Kramosil-Michalek [12]. Latter on Bag and Samanta [2] modified the definition of fuzzy norm of Cheng-Moderson [4] and established the concept of continuity and boundedness of a function with respect to their fuzzy norm in [3].

In this paper, various types of fuzzy anti-continuities and fuzzy anti-boundedness; namely, fuzzy anti-continuity, sequential fuzzy anti-continuity, strong fuzzy anti-continuity, weak fuzzy anti-continuity, strong fuzzy anti-boundedness and weak fuzzy anti-boundedness are defined. The intra relations among fuzzy anti-continuities and intra relation among strongly fuzzy anti-bounded and weakly fuzzy anti-bounded are studied. Also, it is established a very simple

criterion for fuzzy anti-continuity; namely, any linear operator between fuzzy anti-normed linear spaces is strongly and weakly fuzzy anti-continuous if and only if it is strongly and weakly fuzzy anti-bounded respectively.

2 Preliminaries

This section contain some basic definition and preliminary results which will be needed in the sequel.

Definition 2.1 [10]. A binary operation $\diamond : [0, 1] \times [0, 1] \longrightarrow [0, 1]$ is a *t-conorm* if $\diamond$ satisfies the following conditions :

- (i) $\diamond$ is commutative and associative ,
- (ii) $a \diamond 0 = a \quad \forall a \in [0, 1]$,
- (iii) $a \diamond b \leq c \diamond d$ whenever $a \leq c$, $b \leq d$ and $a, b, c, d \in [0, 1]$.

Remark 2.2 [11]. (a) For any $r_1, r_2 \in (0, 1)$ with $r_1 > r_2$, there exist $r_3 \in (0, 1)$ such that $r_1 > r_3 \diamond r_2$.

(b) For any $r_4 \in (0, 1)$, there exist $r_5 \in (0, 1)$ such that $r_5 \diamond r_5 \leq r_4$.

Definition 2.3 [5] Let V be linear space over the field $F (= \mathbf{R} \text{ or } \mathbf{C})$. A fuzzy subset ν of $V \times \mathbf{R}$ is called a **fuzzy antinorm** on V if and only if for all $x, y \in V$ and $c \in F$

- (i) For all $t \in \mathbf{R}$ with $t \leq 0$, $\nu(x, t) = 1$;
- (ii) For all $t \in \mathbf{R}$ with $t > 0$, $\nu(x, t) = 0$ if and only if $x = \theta$;
- (iii) For all $t \in \mathbf{R}$ with $t > 0$, $\nu(cx, t) = \nu(x, \frac{t}{|c|})$ if $c \neq 0, c \in F$;
- (iv) For all $s, t \in \mathbf{R}$ with $\nu(x + y, s + t) \leq \nu(x, s) \diamond \nu(y, t)$;
- (v) $\lim_{t \rightarrow \infty} \nu(x, t) = 0$.

We further assume that for any fuzzy anti-normed linear space (V, A^*) ,

- (vi) $\nu(x, t) < 1, \forall t > 0 \Rightarrow x = \theta$.
- (vii) $\nu(x, \cdot)$ is a continuous function of $\mathbf{R}$ and strictly decreasing on the subset $\{t : 0 < \nu(x, t) < 1\}$ of $\mathbf{R}$.

Theorem 2.4 [5] Let (V, A^*) be a fuzzy antinormed linear space satisfying (vi) and (vii). Let $\|x\|_\alpha^* = \wedge \{t : \nu(x, t) \leq 1 - \alpha\}$, $\alpha \in (0, 1)$. Also, let $\nu' : V \times \mathbf{R} \longrightarrow [0, 1]$ be defined by

$$\begin{aligned} \nu'(x, t) &= \wedge \{1 - \alpha : \|x\|_\alpha^* \leq t\} , \text{ if } (x, t) \neq (\theta, 0) \\ &= 1 , \text{ if } (x, t) = (\theta, 0) \end{aligned}$$

Then $\nu' = \nu$.

Definition 2.5 [1]. Let (U, N^*) be a fuzzy antinormed linear space. A sequence $\{x_n\}_n$ in U is said to be **convergent** to $x \in U$ if given $t > 0$, $r \in (0, 1)$ there exist an integer $n_0 \in \mathbf{N}$ such that

$$N^*(x_n - x, t) < r \quad \forall n \geq n_0.$$

Definition 2.6 [1]. Let (U, N^*) be a fuzzy antinormed linear space. A sequence $\{x_n\}_n$ in U is said to be **cauchy sequence** to $x \in U$ if given $t > 0$, $r \in (0, 1)$ there exist an integer $n_0 \in \mathbf{N}$ such that

$$N^*(x_{n+p} - x_n, t) < r \quad \forall n \geq n_0, p = 1, 2, 3, \dots.$$

Definition 2.7 [1]. A subset A of a fuzzy antinormed linear space (U, N^*) is said to be **bounded** if and only if there exist $t > 0$, $r \in (0, 1)$ such that

$$N^*(x, t) < r \quad \forall x \in A$$

3 Fuzzy Anti-Continuity

Let (U, A^*) and (V, B^*) be any two fuzzy anti-normed linear spaces over the same field F .

Definition 3.1 A mapping $T : (U, A^*) \longrightarrow (V, B^*)$ is said to be **fuzzy anti-continuous** at $x_0 \in U$, if for any given $\epsilon > 0$, $\alpha \in (0, 1)$ there exist $\delta = \delta(\alpha, \epsilon) > 0$, $\beta = \beta(\alpha, \epsilon) \in (0, 1)$ such that for all $x \in U$

$$\nu_U(x - x_0, \delta) < \beta \Rightarrow \nu_V(T(x) - T(x_0), \epsilon) < \alpha.$$

Definition 3.2 A mapping $T : (U, A^*) \longrightarrow (V, B^*)$ is said to be **sequentially fuzzy anti-continuous** at $x_0 \in U$, if for any sequence $\{x_n\}_n$, $x_n \in U$, $\forall n$ with $x_n \longrightarrow x_0$ implies $T(x_n) \longrightarrow T(x_0)$ in V , that is for all $t > 0$,

$$\lim_{n \rightarrow \infty} \nu_U(x_n - x_0, t) = 0 \Rightarrow \lim_{n \rightarrow \infty} \nu_V(T(x_n) - T(x_0), t) = 0.$$

Definition 3.3 A mapping $T : (U, A^*) \longrightarrow (V, B^*)$ is said to be **strongly fuzzy anti-continuous** at $x_0 \in U$, if for any given $\epsilon > 0$ there exist $\delta = \delta(\alpha, \epsilon) > 0$ such that for all $x \in U$,

$$\nu_V(T(x) - T(x_0), \epsilon) \leq \nu_U(x - x_0, \delta)$$

Definition 3.4 A mapping $T : (U, A^*) \longrightarrow (V, B^*)$ is said to be **weakly fuzzy anti-continuous** at $x_0 \in U$, if for any given $\epsilon > 0$, $\alpha \in (0, 1)$ there exist $\delta = \delta(\alpha, \epsilon) > 0$ such that for all $x \in U$,

$$\nu_U(x - x_0, \delta) \leq 1 - \alpha \Rightarrow \nu_V(T(x) - T(x_0), \epsilon) \leq 1 - \alpha.$$

Theorem 3.5 If a mapping T from a fuzzy anti-normed linear space (U, A^*) to a fuzzy anti-normed linear space (V, B^*) is strongly fuzzy anti-continuous then it is weakly fuzzy anti-continuous. But not conversely.

Proof. Obvious.

To show the converse result may not be true we consider the following example.

Example 3.6 As in the example of Note 3.3 of [6], we consider the fuzzy anti-normed linear spaces (X, ν_1) and (X, ν_2) . Let $f(x) = \frac{x^4}{1+x^4} \forall x \in \mathbf{R}$. Now from Example 3 of [8] it directly follows that f is not strongly fuzzy anti-continuous. Here we now show that f is weakly fuzzy anti-continuous on X .

Let $x_0 \in X$, $\epsilon > 0$ and $\delta \in (0, 1)$. Now

$$\nu_2(f(x) - f(x_0), \epsilon) < 1 - \alpha \quad \text{if} \quad \frac{k|f(x) - f(x_0)|}{\epsilon + k|f(x) - f(x_0)|} < 1 - \alpha$$

i.e., if

$$\frac{\epsilon}{\epsilon + k \left| \frac{x^4}{1+x^2} - \frac{x_0^4}{1+x_0^2} \right|} \geq \alpha$$

i.e., if

$$\frac{\frac{\epsilon(1+x^2)(1+x_0^2)}{k|x+x_0||x^2x_0^2+x^2+x_0^2|}}{\frac{\epsilon(1+x^2)(1+x_0^2)}{k|x+x_0||x^2x_0^2+x^2+x_0^2|} + |x-x_0|} \geq \alpha$$

i.e., if

$$\begin{aligned} \alpha |x - x_0| &\leq (1 - \alpha) \frac{\epsilon}{k} \frac{(1+x^2)(1+x_0^2)}{|x+x_0||x^2x_0^2+x^2+x_0^2|} \\ &\leq (1 - \alpha) \frac{\epsilon}{k} \end{aligned}$$

So, depending upon $(1 - \alpha) \frac{\epsilon}{k}$ we may choose $\delta > 0$ such that

$$\alpha(\delta + |x - x_0|) \leq \delta \quad \text{i.e.,} \quad \nu_1(x - x_0, \delta) < 1 - \alpha.$$

Thus we see that for every $\epsilon > 0$, $\alpha \in (0, 1) \exists \delta > 0$ such that

$$\nu_1(x - x_0, \delta) < 1 - \alpha \Rightarrow \nu_2(f(x) - f(x_0), \epsilon) < 1 - \alpha.$$

i.e., f is weakly fuzzy anti-continuous at x_0 .

Theorem 3.7 A mapping T from a fuzzy anti-normed linear space (U, A^*) to a fuzzy anti-normed linear space (V, B^*) is fuzzy anti-continuous if and only if it is sequentially fuzzy anti-continuous.

Proof. The proof of the above theorem is directly follows from Theorem 13 of [8].

Theorem 3.8 *If a mapping T from a fuzzy anti-normed linear space (U, A^*) to a fuzzy anti-normed linear space (V, B^*) is strongly fuzzy anti-continuous then it is sequentially fuzzy anti-continuous.*

Proof. The proof of the above theorem is directly follows from Theorem 12 of [8].

Theorem 3.9 *Let $T : (U, A^*) \longrightarrow (V, B^*)$ be a linear operator. If T is sequentially fuzzy anti-continuous at a point $x_0 \in U$, then it is sequentially fuzzy anti-continuous on U .*

Proof. Let $x \in U$ be an arbitrary point and let $\{x_n\}_n$ be a sequence in U such that $x_n \longrightarrow x$. Then $\forall t > 0$

$$\lim_{n \rightarrow \infty} \nu_U(x_n - x, t) = 0$$

$$\text{i.e., } \lim_{n \rightarrow \infty} \nu_U((x_n - x + x_0) - x_0, t) = 0$$

Since T is sequentially fuzzy anti-continuous at $x_0 \forall t > 0$ we have

$$\lim_{n \rightarrow \infty} \nu_V((x_n - x + x_0) - x_0, t) = 0$$

$$\text{i.e., } \lim_{n \rightarrow \infty} \nu_V(T(x_n) - T(x) + T(x_0) - T(x_0), t) = 0$$

$$\text{i.e., } \lim_{n \rightarrow \infty} \nu_V(T(x_n) - T(x), t) = 0$$

Thus,

$$\lim_{n \rightarrow \infty} \nu_U(x_n - x, t) = 0, \forall t > 0 \Rightarrow \lim_{n \rightarrow \infty} \nu_V(T(x_n) - T(x), t) = 0, \forall t > 0.$$

Hence the proof.

4 Fuzzy Anti-Boundedness

Definition 4.1 *A mapping $T : (U, A^*) \longrightarrow (V, B^*)$ is said to be **strongly fuzzy anti-bounded** on U if and only if there exist a positive real number M such that for all $x \in U$ and for all $t \in R^+$,*

$$\nu_V(T(x), t) \leq \nu_U(x, \frac{t}{M})$$

Example 4.2 *The zero and identity operators are strongly fuzzy anti-bounded.*

Example 4.3 *It is an example of a strongly fuzzy anti-bounded linear operator other than the zero and identity operator.*
Let $(V, \|\cdot\|)$ be a normed linear space over the field $K (= R \text{ or } C)$. Let $\alpha_1, \alpha_2 \in R$ such that $\alpha_1 > \alpha_2 > 0$. Again, let $\nu_1, \nu_2 : V \times R^+ \rightarrow [0, 1]$ be defined by

$$\nu_1(x, t) = \frac{\alpha_1 \|x\|}{t + \alpha_1 \|x\|} \text{ and } \nu_2(x, t) = \frac{\alpha_2 \|x\|}{t + \alpha_2 \|x\|}$$

Also, define $a \diamond b = \max\{a, b\}$ for all $a, b \in [0, 1]$.

Now we shall first show that (V, ν_1) and (V, ν_2) are fuzzy anti-normed linear space.

(i) *The condition (i) is obvious.*

(ii) $\nu_1(x, t) = 0 \Leftrightarrow \frac{\alpha_1 \|x\|}{t + \alpha_1 \|x\|} = 0 \Leftrightarrow \|x\| = 0 \Leftrightarrow x = \theta$.

(iii) *Let $c \in K$ and $c \neq 0$*

$$\begin{aligned} \nu_1(cx, t) &= \frac{\alpha_1 \|cx\|}{t + \alpha_1 \|cx\|} \\ &= \frac{\alpha_1 \|x\|}{\frac{t}{|c|} + \alpha_1 \|x\|} = \nu_1(x, \frac{t}{|c|}) \end{aligned}$$

(iv)

$$\begin{aligned} \nu_1(x + y, s + t) &= \frac{\alpha_1 \|x + y\|}{s + t + \alpha_1 \|x + y\|} \\ &= \frac{1}{\frac{s+t}{\alpha_1 \|x+y\|} + 1} \\ &\leq \frac{1}{\frac{s+t}{\alpha_1 \|x\| + \alpha_1 \|y\|} + 1} \\ &= \frac{\alpha_1 \|x\| + \alpha_1 \|y\|}{s + t + \alpha_1 \|x\| + \alpha_1 \|y\|} \end{aligned}$$

Now if

$$\begin{aligned} \nu_1(x, s) \geq \nu_1(y, t) &\Rightarrow \frac{\alpha_1 \|x\|}{s + \alpha_1 \|x\|} \geq \frac{\alpha_1 \|y\|}{t + \alpha_1 \|y\|}; \\ &\Rightarrow t \|x\| - s \|y\| \end{aligned}$$

Therefore,

$$\frac{\alpha_1 \|x\| + \alpha_1 \|y\|}{s + t + \alpha_1 \|x\| + \alpha_1 \|y\|} - \frac{\alpha_1 \|x\|}{s + \alpha_1 \|x\|} \leq 0.$$

Thus

$$\begin{aligned}\nu_1(x+y, s+t) &\leq \frac{\alpha_1 \|x\| + \alpha_1 \|y\|}{s+t + \alpha_1 \|x\| + \alpha_1 \|y\|} \\ &\leq \frac{\alpha_1 \|x\|}{s + \alpha_1 \|x\|} = \nu_1(x, s) \diamond \nu_1(y, t)\end{aligned}$$

Again if $\nu_1(y, t) \geq \nu_1(x, s)$ Similarly it can be shown that

$$\nu_1(x+y, s+t) \leq \frac{\alpha_1 \|y\|}{t + \alpha_1 \|y\|} = \nu_1(x, s) \diamond \nu_1(y, t)$$

Hence

$$\nu_1(x+y, s+t) \leq \nu_1(x, s) \diamond \nu_1(y, t)$$

(v)

$$\lim_{t \rightarrow \infty} \nu_1(x, t) = \lim_{t \rightarrow \infty} \frac{\alpha_1 \|x\|}{t + \alpha_1 \|x\|} = 0$$

Hence (V, ν_1) is a fuzzy anti-normed linear space. Similarly (V, ν_2) is also fuzzy anti-normed linear space.

We now define a mapping $T : (V, \nu_1) \rightarrow (V, \nu_2)$ by $T(x) = rx$ where $r (\neq 0) \in \mathbf{R}$ is fixed. Clearly T is a linear operator.

Let us choose an arbitrary but fixed $M > 0$ such that $M \geq |r|$ and $x \in V$. Now,

$$\begin{aligned}M \geq |r| &\Rightarrow \alpha_1 M \|x\| \geq \alpha_2 |r| \|x\| \\ &\Rightarrow t + \alpha_1 M \|x\| \geq t + \alpha_2 |r| \|x\| \quad \forall t > 0. \\ &\Rightarrow \frac{t}{t + \alpha_2 |r| \|x\|} \geq \frac{t}{t + \alpha_1 M \|x\|} \quad \forall t > 0. \\ &\Rightarrow \frac{t}{t + \alpha_2 \|rx\|} \geq \frac{\frac{t}{M}}{\frac{t}{M} + \alpha_1 \|x\|} \quad \forall t > 0. \\ &\Rightarrow 1 - \frac{t}{t + \alpha_2 \|rx\|} \leq 1 - \frac{\frac{t}{M}}{\frac{t}{M} + \alpha_1 \|x\|} \quad \forall t > 0. \\ &\Rightarrow \frac{\alpha_2 \|rx\|}{t + \alpha_2 \|rx\|} \leq \frac{\alpha_1 \|x\|}{\frac{t}{M} + \alpha_1 \|x\|} \quad \forall t > 0.\end{aligned}$$

i.e.,

$$\nu_2(T(x), t) \leq \nu_1(x, \frac{t}{M}) \quad \forall t > 0 \text{ and } \forall x \in V.$$

Hence T is strongly fuzzy anti-bounded on V .

Definition 4.4 A mapping $T : (U, A^*) \longrightarrow (V, B^*)$ is said to be **weakly fuzzy anti-bounded** on U if and only if for any $\alpha \in (0, 1)$ there exist $M_\alpha (> 0)$ such that for all $x \in U$ and for all $t \in R^+$,

$$\nu_U(x, \frac{t}{M_\alpha}) \leq 1 - \alpha \Rightarrow \nu_V(T(x), t) \leq 1 - \alpha$$

Theorem 4.5 Let $T : (U, A^*) \longrightarrow (V, B^*)$ be a linear operator. If T is strongly fuzzy anti-bounded then it is weakly fuzzy anti-bounded. But not conversely.

Proof. First we suppose that T is strongly fuzzy anti-bounded. Then there exist $M > 0$ such that $\forall x \in U$ and $\forall t \in R$,

$$\nu_V(T(x), t) \leq \nu_U(x, \frac{t}{M})$$

Thus for any $\alpha \in (0, 1)$, there exists $M_\alpha (= M)$ such that

$$\nu_U(x, \frac{t}{M_\alpha}) \leq 1 - \alpha \Rightarrow \nu_V(T(x), t) \leq 1 - \alpha$$

Hence T is weakly fuzzy anti-bounded.

The converse of the above theorem is not necessarily true. For example

Example 4.6 Let $(V, \|\cdot\|)$ be a linear space over the field $K (= R \text{ or } C)$ and $\nu_1, \nu_2 : V \times R \longrightarrow [0, 1]$ be defined by

$$\begin{aligned} \nu_1(x, t) &= \frac{2\|x\|^2}{t^2 + \|x\|^2} \quad , \text{ if } t > \|x\| \\ &= 1 \quad , \text{ if } 0 < t \leq \|x\| \end{aligned}$$

and
$$\nu_2(x, t) = \frac{\|x\|}{t + \|x\|}$$

Also define $a \diamond b = \max\{a, b\}$

Already we have seen that (V, ν_2) is a fuzzy anti-normed linear space. Now we shall prove that (V, ν_1) is a fuzzy anti-normed linear space.

(i) The condition (i) is obvious.

(ii) $\nu_1(x, t) = 0 \Leftrightarrow \frac{2\|x\|^2}{t^2 + \|x\|^2} = 0 \Leftrightarrow \|x\| = 0 \Leftrightarrow x = \theta$.

(iii) Let, $c \in K$ and $c \neq 0$. If $t > \|cx\|$,

$$\nu_1(cx, t) = \frac{2\|cx\|^2}{t^2 + \|cx\|^2} = \frac{2|c|^2\|x\|^2}{t^2 + |c|^2\|x\|^2} = \frac{2\|x\|^2}{(\frac{t}{|c|})^2 + \|x\|^2} = \nu_1(x, \frac{t}{|c|})$$

Again if $0 < t \leq \|cx\|$ then $\nu_1(cx, t) = 1$

and $0 < t \leq \|cx\| \Rightarrow 0 < \frac{t}{|c|} \leq \|x\| \Rightarrow \nu_1(x, \frac{t}{|c|}) = 1$

(iv) Let $s, t \in \mathbf{R}^+$, $x, y \in V$

If $0 < s + t \leq \|x + y\|$, we have the following possibilities

$$(a) 0 < s \leq \|x\| \text{ and } 0 < t \leq \|y\|$$

$$(b) 0 < s \leq \|x\| \text{ and } t > \|y\|$$

$$(c) 0 < t \leq \|y\| \text{ and } s > \|x\|$$

In each case $\nu_1(x + y, s + t) = 1 = \nu_1(x, s) \diamond \nu_1(y, t)$ Again, if $s + t > \|x + y\|$, we have the following four possibilities

$$(a) s > \|x\|, t \leq \|y\|$$

$$(b) s \leq \|x\|, t > \|y\|$$

$$(c) s \leq \|x\|, t \leq \|y\|$$

$$(d) s > \|x\|, t > \|y\|$$

In the cases (a), (b), (c)

$$\begin{aligned} \nu_1(x + y, s + t) &= \frac{2\|x + y\|^2}{(s + t)^2 + \|x + y\|^2} \\ &< 1 = \nu_1(x, s) \diamond \nu_1(y, t) \end{aligned}$$

So, we now suppose that $s > \|x\|$ and $t > \|y\|$ Now, $s + t > \|x\| + \|y\| \geq \|x + y\|$. Therefore,

$$\begin{aligned} \nu_1(x + y, s + t) &= \frac{2\|x + y\|^2}{(s + t)^2 + \|x + y\|^2} \\ &\leq \frac{2(\|x\| + \|y\|)^2}{(s + t)^2 + (\|x\| + \|y\|)^2} \end{aligned}$$

Hence we have

$$\begin{aligned} \nu_1(x + y, s + t) &\leq \frac{2(\|x\| + \|y\|)^2}{(s + t)^2 + (\|x\| + \|y\|)^2} \\ &\leq \frac{2\|y\|^2}{t^2 + \|y\|^2} = \nu_1(y, t) \end{aligned}$$

when $\nu_1(x, s) \leq \nu_1(y, t)$

Similarly,

$$\nu_1(x + y, s + t) \leq \frac{2(\|x\| + \|y\|)^2}{(s + t)^2 + (\|x\| + \|y\|)^2}$$

$$\leq \frac{2\|x\|^2}{t^2 + \|x\|^2} = \nu_1(x, s)$$

when $\nu_1(y, t) \leq \nu_1(x, s)$

Thus

$$\nu_1(x + y, s + t) \leq \nu_1(x, s) \diamond \nu_1(y, t)$$

$$(v) \lim_{t \rightarrow \infty} \nu_1(x, t) = \lim_{t \rightarrow \infty} \frac{2\|x\|^2}{t^2 + \|x\|^2} = 0$$

Thus we see that (V, ν_1) is a fuzzy anti-normed linear space.

Now we define a linear operator $T : (U, \nu_1) \longrightarrow (V, \nu_2)$ by $T(x) = x$, $\forall x \in V$. Let, $\alpha \in (0, 1)$, $x \in V$ and $t \in R^+$ and choose $M_\alpha = \frac{1}{1-\alpha}$. We now prove that

$$\nu_1(x, \frac{t}{M_\alpha}) \leq 1 - \alpha \Rightarrow \nu_2(T(x), t) \leq 1 - \alpha.$$

$$\begin{aligned} \nu_1(x, \frac{t}{M_\alpha}) &\leq 1 - \alpha \\ \Rightarrow \frac{2\|x\|^2}{t^2(1-\alpha)^2 + \|x\|^2} &\leq 1 - \alpha \\ \Rightarrow 1 - \frac{2\|x\|^2}{t^2(1-\alpha)^2 + \|x\|^2} &\geq 1 - (1 - \alpha) = \alpha \\ \Rightarrow \frac{t^2(1-\alpha)^2 - \|x\|^2}{t^2(1-\alpha)^2 + \|x\|^2} &\geq \alpha \\ \Rightarrow t^2(1-\alpha)^3 &\geq (1+\alpha)\|x\|^2 \\ \Rightarrow \|x\| &\leq \frac{t(1-\alpha)\sqrt{1-\alpha}}{\sqrt{1+\alpha}} \\ \Rightarrow t + \|x\| &\leq t \frac{(1-\alpha)\sqrt{1-\alpha} + \sqrt{1+\alpha}}{\sqrt{1+\alpha}} \\ \Rightarrow \frac{t}{t + \|x\|} &\geq \frac{\sqrt{1+\alpha}}{(1-\alpha)\sqrt{1-\alpha} + \sqrt{1+\alpha}} \\ \Rightarrow 1 - \frac{t}{t + \|x\|} &\leq 1 - \frac{\sqrt{1+\alpha}}{(1-\alpha)\sqrt{1-\alpha} + \sqrt{1+\alpha}} \\ \Rightarrow \frac{\|x\|}{t + \|x\|} &\leq \frac{(1-\alpha)\sqrt{1-\alpha}}{(1-\alpha)\sqrt{1-\alpha} + \sqrt{1+\alpha}} \end{aligned}$$

Now

$$\begin{aligned} \frac{(1-\alpha)\sqrt{1-\alpha}}{(1-\alpha)\sqrt{1-\alpha} + \sqrt{1+\alpha}} &\leq 1-\alpha \Leftrightarrow \sqrt{1-\alpha} \leq (1-\alpha)\sqrt{1-\alpha} + \sqrt{1+\alpha} \\ &\Leftrightarrow \alpha\sqrt{1-\alpha} \leq \sqrt{1+\alpha} \\ &\Leftrightarrow 1 + \alpha + \alpha^3 \geq \alpha^2 \end{aligned}$$

which is true for all $\alpha \in (0, 1)$.

Hence

$$\nu_1\left(x, \frac{t}{M_\alpha}\right) \leq 1-\alpha \Rightarrow \nu_2(T(x), t) \leq 1-\alpha.$$

Thus T is weakly fuzzy anti-bounded on X .

Definition 4.7 A linear operator $T : (U, A^*) \longrightarrow (V, B^*)$ is said to be **uniformly fuzzy anti-bounded** if and only if there exist $M > 0$ such that

$$\|T(x)\|_\alpha^* \geq M \|x\|_\alpha^*, \quad \alpha \in (0, 1)$$

where $\{\|\cdot\|_\alpha^* : \alpha \in (0, 1)\}$ is ascending family of fuzzy α -anti-norms.

Theorem 4.8 Let $T : (U, A^*) \longrightarrow (V, B^*)$ be a linear operator and (U, A^*) and (V, B^*) satisfies (vi) and (vii). Then T is strongly fuzzy anti-bounded if and only if it is uniformly fuzzy anti-bounded with respect to fuzzy α -anti-norms.

Proof. Let $\{\|\cdot\|_\alpha^* : \alpha \in (0, 1)\}$ be ascending family of α -norms. First suppose that T is strongly fuzzy anti-bounded. Then there exist $M > 0$ such that $\forall x \in U$ and $\forall s \in R$,

$$\nu_V(T(x), t) \leq \nu_U\left(x, \frac{s}{M}\right) \text{ i.e., } \nu_V(T(x), t) \leq \nu_U(Mx, s)$$

$$\|Mx\|_\alpha^* > t \Rightarrow \wedge \{s : \nu(Mx, s) \leq 1-\alpha\} > t.$$

$$\Rightarrow \exists s_0 > t \text{ such that } \nu(Mx, s_0) \leq 1-\alpha$$

$$\Rightarrow \exists s_0 > t \text{ such that } \nu(T(x), s_0) \leq 1-\alpha$$

$$\Rightarrow \|T(x)\|_\alpha^* \geq s_0 > t$$

Hence $\|T(x)\|_\alpha^* \geq \|Mx\|_\alpha^* = M \|x\|_\alpha^*$.

Thus T is uniformly fuzzy anti-bounded.

Conversely, suppose that there exist $M > 0$ such that $\forall x \in U$ and $\forall \alpha \in (0, 1)$

$$\|T(x)\|_\alpha^* \geq M \|x\|_\alpha^*$$

Let, $p > \nu_U(Mx, s) \Rightarrow p > \wedge \{ \alpha \in (0, 1) : \| Mx \|_\alpha^* \leq s \}$
 $\Rightarrow$ There exist $\alpha_0 \in (0, 1)$ such that $p > \alpha_0$ and $\| Mx \|_\alpha^* \leq s$
 $\Rightarrow \| T(x) \|_\alpha^* \leq s$
 $\Rightarrow \nu_V(T(x), s) \leq 1 - \alpha_0 < p$.
Hence, $\nu_V(T(x), s) \leq \nu_U(Mx, s) = \nu_U(x, \frac{s}{M})$.
Thus T is strongly fuzzy anti-bounded.

Theorem 4.9 Let $T : (U, A^*) \longrightarrow (V, B^*)$ be a linear operator. Then,

- (i) T is strongly fuzzy anti-continuous on U if T is strongly fuzzy anti-continuous at a point $x_0 \in U$.
- (ii) T is strongly fuzzy anti-continuous if and only if T is strongly fuzzy anti-bounded.

Proof. (i) Since, T is strongly fuzzy anti-continuous at $x_0 \in U$, for each $\epsilon > 0$ there exists $\delta > 0$ such that

$$\nu_V(T(x) - T(x_0), \epsilon) \leq \nu_U(x - x_0, \delta)$$

Taking $y \in U$ and replacing x by $x + x_0 - y$, we get,

$$\begin{aligned} \nu_V(T(x) - T(x_0), \epsilon) &\leq \nu_U(x - x_0, \delta) \\ \Rightarrow \nu_V(T(x + x_0 - y) - T(x_0), \epsilon) &\leq \nu_U(x + x_0 - y - x_0, \delta) \\ \Rightarrow \nu_V(T(x) + T(x_0) - T(y) - T(x_0), \epsilon) &\leq \nu_U(x - y, \delta) \\ \Rightarrow \nu_V(T(x) - T(y), \epsilon) &\leq \nu_U(x - y, \delta) \end{aligned}$$

Since, $y \in U$ is arbitrary, T is strongly fuzzy anti-continuous on U .

(ii) First we suppose that T is strongly fuzzy anti-bounded. Thus there exist a positive real number M such that for all $x \in U$ and for all $\epsilon \in R^+$,

$$\nu_V(T(x), \epsilon) \leq \nu_U(x, \frac{\epsilon}{M})$$

$$i.e., \nu_V(T(x) - T(\theta), \epsilon) \leq \nu_U(x - \theta, \frac{\epsilon}{M})$$

$$i.e., \nu_V(T(x) - T(\theta), \epsilon) \leq \nu_U(x - \theta, \delta)$$

where $\delta = \frac{\epsilon}{M}$.

Thus T is strongly fuzzy anti-continuous at θ and hence T is strongly fuzzy anti-continuous on U .

Conversely, suppose that T is strongly fuzzy anti-continuous on U . Using fuzzy anti-continuity of T at $x = \theta$ for $\epsilon = 1$ there exist $\delta > 0$ such that for all $x \in U$,

$$\nu_V(T(x) - T(\theta), 1) \leq \nu_U(x - \theta, \delta)$$

If $x \neq \theta$ and $t > 0$. Putting $x = ut$

$$\nu_V(T(x), t) = \nu_V(ut, t) = \nu_V(T(u), 1) \leq \nu_U(u, \delta) = \nu_U\left(\frac{x}{t}, \delta\right) = \nu_U\left(x, \frac{t}{M}\right), \text{ Where } M = \frac{1}{\delta}.$$

So, $\nu_V(T(x), t) \leq \nu_U\left(x, \frac{t}{M}\right)$

If $x \neq \theta$ and $t \leq 0$ then $\nu_V(T(x), t) = 1 = \nu_U\left(x, \frac{t}{M}\right)$.

If $x = \theta$ and $t \in R$, then $T(\theta_U) = \theta_V$ and

$$\nu_V(\theta_V, t) = \nu_U\left(\theta_U, \frac{t}{M}\right) = 0, \text{ if } t > 0.$$

$$\nu_V(\theta_V, t) = \nu_U\left(\theta_U, \frac{t}{M}\right) = 1, \text{ if } t \leq 0.$$

Hence T is strongly fuzzy anti-bounded.

Theorem 4.10 Let $T : (U, A^*) \longrightarrow (V, B^*)$ be a linear operator. Then,

(i) T is weakly fuzzy anti-continuous on U if T is weakly fuzzy anti-continuous at a point $x_0 \in U$.

(ii) T is weakly fuzzy anti-continuous if and only if T is weakly fuzzy anti-bounded.

Proof. (i) Since, T is weakly fuzzy anti-continuous at x_0 in U , for $\epsilon > 0$ and $\alpha \in (0, 1)$ there exist $\delta = \delta(\alpha, \epsilon) > 0$ such that $\forall x \in U$

$$\nu_U(x - x_0, \delta) \leq 1 - \alpha \Rightarrow \nu_V(T(x) - T(x_0), \epsilon) \leq 1 - \alpha.$$

Taking $y \in U$ and replacing x by $x + x_0 - y$ we get,

$$\nu_U(x + x_0 - y - x_0, \delta) \leq 1 - \alpha \Rightarrow \nu_V(T(x + x_0 - y) - T(x_0), \epsilon) \leq 1 - \alpha$$

$$\text{i.e., } \nu_U(x - y, \delta) \leq 1 - \alpha \Rightarrow \nu_V(T(x) + T(x_0) - T(y) - T(x_0), \epsilon) \leq 1 - \alpha$$

$$\text{i.e., } \nu_U(x - y, \delta) \leq 1 - \alpha \Rightarrow \nu_V(T(x) - T(y), \epsilon) \leq 1 - \alpha$$

Since, $y (\in U)$ is arbitrary it follows that T is weakly fuzzy anti-continuous on U .

(ii) First we suppose that T is fuzzy anti-bounded. Thus for any $\alpha \in (0, 1)$ there exist $M_\alpha > 0$ such that $\forall t \in R$ and $\forall x \in U$ we have

$$\nu_U\left(x, \frac{t}{M}\right) \leq 1 - \alpha \Rightarrow \nu_V(T(x), t) \leq 1 - \alpha$$

Therefore,

$$\nu_U\left(x - \theta, \frac{t}{M}\right) \leq 1 - \alpha \Rightarrow \nu_V(T(x) - T(\theta), t) \leq 1 - \alpha$$

$$i.e., \nu_U(x-\theta, \frac{\epsilon}{M_\alpha}) \leq 1-\alpha \Rightarrow \nu_V(T(x)-T(\theta), \epsilon) \leq 1-\alpha$$

$$i.e., \nu_U(x-\theta, \delta) \leq 1-\alpha \Rightarrow \nu_V(T(x)-T(\theta), \epsilon) \leq 1-\alpha$$

where $\delta = \frac{\epsilon}{M_\alpha}$

Thus, T is weakly fuzzy anti-continuous at x_0 and hence weakly fuzzy anti-continuous on U .

Conversely, suppose that T is weakly fuzzy anti-continuous on U . Using continuity of T at θ and taking $\epsilon = 1$ we have for all $\alpha \in (0, 1)$ there exists $\delta(\alpha, 1) > 0$ such that for all $x \in U$,

$$\nu_U(x - \theta, \delta) \leq 1 - \alpha \Rightarrow \nu_V(T(x) - T(\theta), 1) \leq 1 - \alpha$$

$$i.e., \nu_U(x, \delta) \leq 1 - \alpha \Rightarrow \nu_V(T(x), 1) \leq 1 - \alpha.$$

If $x \neq \theta$ and $t > 0$. Putting $x = \frac{u}{t}$ we have,

$$\nu_U(\frac{u}{t}, \delta) \leq 1 - \alpha \Rightarrow \nu_V(T(\frac{u}{t}), 1) \leq 1 - \alpha$$

$$i.e., \nu_U(u, t\delta) \leq 1 - \alpha \Rightarrow \nu_V(T(u), t) \leq 1 - \alpha$$

$$i.e., \nu_U(u, \frac{t}{M_\alpha}) \leq 1 - \alpha \Rightarrow \nu_V(T(\frac{u}{t}), 1) \leq 1 - \alpha$$

where $M_\alpha = \frac{1}{\delta(\alpha, 1)}$ If $x \neq \theta$ and $t \leq 0$, $\nu_U(x, \frac{t}{M_\alpha}) = \nu_V(T(x), t) = 1$ for any $M_\alpha > 0$.

If $x = \theta$ then for $M_\alpha > 0$,

$$\nu_U(x, \frac{t}{M_\alpha}) = \nu_V(T(x), t) = 0 \quad , \text{ if } t > 0$$

$$\nu_U(x, \frac{t}{M_\alpha}) = \nu_V(T(x), t) = 1 \quad , \text{ if } t \leq 0$$

Hence, T is weakly fuzzy anti-bounded.

Theorem 4.11 *Let $T : (U, A^*) \longrightarrow (V, B^*)$ be a linear operator and (U, A^*) and (V, B^*) satisfies (vi) and (vii). Then T is weakly fuzzy anti-bounded if and only if T is fuzzy anti-bounded with respect to α -norms.*

Proof. First we suppose that T is weakly fuzzy anti-bounded. Then for all $\alpha \in (0, 1)$ there exist $M_\alpha > 0$ such that $\forall x \in U, t \in R$ we have

$$\nu_U(x, \frac{t}{M_\alpha}) \leq 1 - \alpha \Rightarrow \nu_V(T(x), t) \leq 1 - \alpha$$

Hence we get, $\nu_U(M_\alpha x, t) \leq 1 - \alpha \Rightarrow \nu_V(T(x), t) \leq 1 - \alpha$

$i.e., \bigwedge \{ \beta \in (0, 1) : \| M_\alpha x \|_\beta^* \leq t \} \leq 1 - \alpha \Rightarrow \bigwedge \{ \beta \in (0, 1) :$

$$\|T(x)\|_{\beta}^* \leq t \} \leq 1 - \alpha$$

Now we show that

$$\wedge \{ \beta \in (0, 1) : \|M_{\alpha}x\|_{\beta}^* \leq t \} \leq 1 - \alpha \Leftrightarrow \|M_{\alpha}x\|_{\alpha}^* \leq t$$

If $x = \theta$ then the relation is obvious.

Suppose $x \neq \theta$.

Now, if

$$\wedge \{ \beta \in (0, 1) : \|M_{\alpha}x\|_{\beta}^* \leq t \} < 1 - \alpha \text{ then } \|M_{\alpha}x\|_{\alpha}^* \leq t \quad (1)$$

If $\wedge \{ \beta \in (0, 1) : \|M_{\alpha}x\|_{\beta}^* \leq t \} = 1 - \alpha$ then there exists a decreasing sequence $\{\alpha_n\}_n$ in $(0, 1)$ such that $\alpha_n \rightarrow \alpha$ and $\|M_{\alpha}x\|_{\alpha}^* \leq t$ Then by Theorem 3.7 we have

$$\|M_{\alpha}x\|_{\alpha}^* \leq t \quad (2)$$

From (1) and (2) we get

$$\wedge \{ \beta \in (0, 1) : \|M_{\alpha}x\|_{\beta}^* \leq t \} \leq 1 - \alpha \Rightarrow \|M_{\alpha}x\|_{\alpha}^* \leq t \quad (3)$$

Next we suppose that $\|M_{\alpha}x\|_{\alpha}^* \leq t$.

If $\|M_{\alpha}x\|_{\alpha}^* < t$ then $\nu_U(M_{\alpha}x, t) \leq 1 - \alpha$. i.e.,

$$\wedge \{ \beta \in (0, 1) : \|M_{\alpha}x\|_{\beta}^* \leq t \} \leq 1 - \alpha \quad (4)$$

If $\|M_{\alpha}x\|_{\alpha}^* = t$ i.e., $\wedge \{ s : \nu_U(M_{\alpha}x, s) \leq 1 - \alpha \} = t$ then there exist an increasing sequence $\{s_n\}_n$ in $\mathbf{R}^+$ such that $s_n \rightarrow t$ and

$$\nu_U(M_{\alpha}x, s_n) \leq 1 - \alpha \Rightarrow \lim_{n \rightarrow \infty} \nu_U(M_{\alpha}x, s_n) \leq 1 - \alpha$$

$$\Rightarrow \nu_U(M_{\alpha}x, \lim_{n \rightarrow \infty} s_n) \leq 1 - \alpha$$

$$\Rightarrow \nu_U(M_{\alpha}x, t) \leq 1 - \alpha$$

$$\Rightarrow \wedge \{ \beta \in (0, 1) : \|M_{\alpha}x\|_{\beta}^* \leq t \} \leq 1 - \alpha$$

Hence from (4) it follows that

$$\|M_{\alpha}x\|_{\alpha}^* \leq t \Rightarrow \wedge \{ \beta \in (0, 1) : \|M_{\alpha}x\|_{\beta}^* \leq t \} \leq 1 - \alpha \quad (5)$$

From (3) and (5) we have

$$\wedge \{ \beta \in (0, 1) : \|M_{\alpha}x\|_{\beta}^* \leq t \} \leq 1 - \alpha \Leftrightarrow \|M_{\alpha}x\|_{\alpha}^* \leq t \quad (6)$$

In the similar way we can show that

$$\wedge \{ \beta \in (0, 1) : \|T(x)\|_{\beta}^* \leq t \} \leq 1 - \alpha \Leftrightarrow \|T(x)\|_{\alpha}^* \leq t \quad (7)$$

From (6) and (7) we have $\nu_U(M_\alpha x, t) \leq 1 - \alpha \Rightarrow \nu_V(T(x), t) \leq 1 - \alpha$
Then

$$\|M_\alpha x\|_\alpha^* \leq t \Rightarrow \|T(x)\|_\alpha^* \leq t$$

This implies that

$$\|T(x)\|_\alpha^* \geq \|M_\alpha x\|_\alpha^*$$

Conversely, suppose that $\forall \alpha \in (0, 1), \exists M_\alpha > 0$ such that $\forall x \in U$,

$$\|T(x)\|_\alpha^* \geq \|M_\alpha x\|_\alpha^*$$

Then for $x \neq \theta$ and $\forall t > 0$,

$$\|M_\alpha x\|_\alpha^* \leq t \Rightarrow \|T(x)\|_\alpha^* \leq t$$

i.e.,

$$\wedge \{s : \nu_U(M_\alpha x, s) \leq 1 - \alpha\} \leq t \Rightarrow \wedge \{s : \nu_V(T(x), s) \leq 1 - \alpha\} \leq t$$

In the similar way as above we can show that

$$\wedge \{s : \nu_U(M_\alpha x, s) \leq 1 - \alpha\} \leq t \Leftrightarrow \nu_U(M_\alpha x, t) \leq 1 - \alpha$$

and

$$\wedge \{s : \nu_U(T(x), s) \leq 1 - \alpha\} \leq t \Leftrightarrow \nu_U(T(x), t) \leq 1 - \alpha$$

Thus we have

$$\nu_U(x, \frac{t}{M_\alpha}) \leq 1 - \alpha \Rightarrow \nu_V(T(x), t) \leq 1 - \alpha, \forall x \in U$$

If $x \neq \theta, t \leq 0$ and if $x = \theta, t > 0$ then the above relation is obvious.
Hence the proof.

Theorem 4.12 *Let, $T : (U, A^*) \longrightarrow (V, B^*)$ be a linear operator and (U, A^*) and (V, B^*) satisfies (vi) and (vii). If U is finite dimensional then T is weakly fuzzy anti-bounded.*

Proof. Since, (U, A^*) and (V, B^*) satisfies (vi), we may suppose that $\{\|\cdot\|_\alpha^* : \alpha \in (0, 1)\}$ is ascending family of fuzzy α -anti-norms.

Since T is of finite dimension, $T : (U, A^*) \longrightarrow (V, B^*)$ is bounded linear operator for each $\alpha \in (0, 1)$. Thus by Theorem 4.11 it follows that T is weakly fuzzy anti-bounded.

5 Future Work

In our next paper we shall try to develop the concept of fuzzy anti-bounded linear functionals and their properties.

References

- [1] Iqbal H. Jebril and T.K. Samanta , *Fuzzy anti-normed linear space*, Journal of mathematics and Technology, February, 2010.
- [2] T. Bag and S.K. Samanta , *Finite Dimensional Fuzzy Normed Linear Spaces*, The Journal of Fuzzy Mathematics Vol. 11 (2003) 687 - 705.
- [3] T. Bag and S.K. Samanta , *Fuzzy bounded linear operators* , Fuzzy Sets and Systems 151 (2005) 513 - 547.
- [4] S.C. Cheng and J.N. Mordeson , *Fuzzy Linear Operators and Fuzzy Normed Linear Spaces* , Bull. Cal. Math. Soc.86 (1994) 429 - 436.
- [5] Bivas Dinda, T.K. Samanta and Iqbal H. Jebril , *Fuzzy Anti-norm and Fuzzy α -anti-convergence* , (Comunicated)
- [6] Bivas Dinda and T.K. Samanta , *Intuitionistic Fuzzy Continuity and Uniform Convergence* , Int. J. Open Problems Compt.Math., Vol 3, No. 1 (2010) 8 - 26.
- [7] C. Felbin , *The completion of fuzzy normed linear space*, Journal of mathematical analysis and application 174(2) (1993) 428-440.
- [8] T.K. Samanta and Iqbal H. Jebril , *Finite dimentional intuitionistic fuzzy normed linear space*, Int. J. Open Problems Compt. Math., Vol 2, No. 4 (2009) 574-591.
- [9] A.K. Katsaras , *Fuzzy topological vector space*, Fuzzy Sets and Systems 12 (1984) 143 - 154.
- [10] B. Schweizer , A. Sklar, *Statistical metric space*, Pacific journal of mathematics 10 (1960) 314-334.
- [11] S. Vijayabalaji, N. Thillaigovindan, Y.B. Jun *Intuitionistic Fuzzy n -normed linear space* , Bull. Korean Math. Soc. 44 (2007) 291 - 308.
- [12] O. Kramosil, J. Michalek , *Fuzzy metric and statisticalmetric spaces*, Kybernetika 11 (1975) 326 - 334.
- [13] L.A. Zadeh *Fuzzy sets*, Information and control 8 (1965) 338-353.